PROSPECTS FOR FUNDAMENTAL SYMMETRY TESTS WITH POLYATOMIC MOLECULES

ROBERT BERGER and TIMUR ISAEV, *Clemens-Schöpf Institute, Technical University of Darmstadt, 64287 Darmstadt, Germany*.

Special features of polyatomic molecules make them attractive candidates for search for violation of fundamental symmetries and variation of fundamental constants [1, 2]. We discuss the possibility of searching for nuclear spin-dependent space-parity violating ($\mathcal{NSD} - \mathcal{PV}$) interaction in closed-shell and open-shell polyatomic molecules. The parameter W_{a} of the effective molecular spin-rotational Hamiltonian characterising the strength of $\mathcal{NSD} - \mathcal{PV}$ interaction in open-shell linear molecules is discussed and approaches for its calculation outlined. In addition, possibilities for detecting $\mathcal{NSD} - \mathcal{PV}$ in chiral molecules via NMR and MW spectroscopy are presented.

REFERENCES:

1. C. Stoeffler et al, Phys. Chem. Chem. Phys., 13 (3), 2011; M. Quack, J. Stohner and M. Willeke, Ann. Rev. Phys. Chem., 59, 2008
2. J. Bagdonaite et al, Science, 339 (6115), 2013.