

OVERTONE AND COMBINATION BAND SPECTROSCOPY OF H_3^+

BENJAMIN J. McCALL^a, *Department of Chemistry and Department of Astronomy and Astrophysics, The University of Chicago, Chicago, IL 60637*; THERESE R. HUET, *Laboratoire de Spectroscopie Hertzienne, Centre d'Etudes et de Recherches Lasers et Applications, Universite de Lille, 59 655 Villeneuve d'Ascq cedex, France*; JAMES K. G. WATSON, *Steacie Institute for Molecular Sciences, National Research Council of Canada, Ottawa, Ontario, Canada K1A 0R6*; TAKESHI OKA, *Department of Chemistry and Department of Astronomy and Astrophysics, The University of Chicago, Chicago, IL 60637*.

We report our detection of H_3^+ rovibrational transitions between 7700 and 8200 cm^{-1} . This work employed a New Focus external cavity tunable diode laser, a positive column discharge, and the velocity modulation technique for enhanced sensitivity to molecular ions. This frequency range encompasses transitions of the $3\nu_2^1 \leftarrow 0$ and $\nu_1 + 2\nu_2 \leftarrow 0$ bands — the highest energy transitions studied to date.

As the simplest three-atom molecule, H_3^+ has long served as a benchmark for *ab initio* calculations. We have compared our observed frequencies with our predictions, and expect that this work (along with detection of even higher overtones) will lead to better *ab initio* treatments of this important molecule.

^aSupported by the Fannie and John Hertz Foundation