MICROWAVE AND *AB INITIO* INVESTIGATION OF $(CH_3)_3N\cdots HF\cdots HF$

C. S. BRAUER, M. B. CRADDOCK, S. W. HUNT, and K. R. LEOPOLD, *Department of Chemistry, University of Minnesota, Minneapolis, MN 55455*; JENS UWE-GRABOW, *Lehrgebiet Physikalische Chemie A, Institut für Physikalische Chemie und Elektrochemie, Universtät Hannover, Callinstr. 3-3a, D-30167 Hannover*.

Rotational spectroscopy and *ab initio* calculations have been used to examine the effect of a single HF solvent molecule on the gas phase proton transfer in the hydrogen bonded complex $(CH_3)_3N\cdots HF$. The rotational spectra of $(CH_3)_3N\cdots HF\cdots HF$ and five of its isotopically substituted derivatives have been observed by Fourier transform microwave spectroscopy. This follows a previous study by our group on $H_3N\cdots HF\cdots HF$, in which the addition of the second HF decreased the $N\cdots H$ hydrogen bond distance in $H_3N\cdots HF$ by 0.21(6) $\mathring{A}$. The present study investigates the effect of the increased basicity of the amine on the $N\cdots H$ hydrogen bond. We observe a simple asymmetric rotor spectrum with strong a- and b- type transitions, consistent with a ring structure for the $N\cdots HF\cdots HF$ frame. No evidence of internal rotation is observed. Structural analysis is underway and will be discussed, but preliminary analysis indicates an $N\cdots H$ hydrogen bond distance of about 1.4 $\mathring{A}$, approximately halfway between the hydrogen bond distance in $H_3N\cdots HF$ of 1.7 $\mathring{A}$ and the N-H covalent length of 1.1 $\mathring{A}$. *Ab initio* calculations concur with experiment, indicating that, as with $H_3N\cdots HF\cdots HF$, the trimethylamine complex forms a ring in which both the $N\cdots H$ hydrogen bond and the HFH angle are significantly perturbed. In addition, two of the methyl groups of the trimethylamine assume an eclipsed conformation, apparently participating in a bifurcated interaction with the fluorine atom of the second HF. This complex provides the first step in microsolvation of $(CH_3)_3N\cdots HF$ and is useful in understanding the role of local environment in promoting proton transfer.