

SUB-DOPPLER HIGH RESOLUTION EXCITATION SPECTROSCOPY OF DIBENZOFURAN

MASAAKI BABA, *Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan*; ATSUSHI DOI, YOSHIO TATAMITANI, MICHIRU YAMAWAKI, SHUNJI KASAHARA, and HAJIME KATÔ, *Molecular Photoscience Research Center, Kobe University, Kobe 657-8501, Japan*.

Dibenzofuran (DF) is one of the prototypical molecules of toxic dioxins as well as dibenzo-*p*-dioxin.^a Sub-Doppler high-resolution fluorescence excitation spectra of the $S_1 \leftarrow S_0$ transition of DF have been observed in a collimated molecular beam. Each rotational line was fully resolved and the rotational constants in both the S_0 and S_1 states were determined by the rotational analysis of the 0-0 band ($33773.478 \text{ cm}^{-1}$). The DF molecule has shown to be planar both in the S_0 and S_1 states. All of the observed lines were the $\Delta K = \pm 1$ transition (B type). Therefore, the S_1 state is assigned to be the 1A_1 state in C_{2v} symmetry.

^aM. Baba, A. Doi, Y. Tatamitani, S. Kasahara, and H. Katô, *J. Phys. Chem. A*, **108**, 1388 (2004)