

STUDY OF THE GLYCOLALDEHYDE AND GLYCOLALDEHYDE-WATER COMPLEX BY AB INITIO CALCULATIONS AND FTMW SPECTROSCOPY IN A SUPERSONIC MOLECULAR BEAM.

J.R. AVILES-MORENO, D. PETITPREZ, and T.R. HUET, *Laboratoire PhLAM, UMR 8523 CNRS-Universite de Lille 1, F-59655 Villeneuve d'Ascq Cedex, France.*

We are interested in studying sugars ($C_nH_{2n}O_n$) because of their biological role. Also, the spectroscopic characterization of their hydrated complexes could help to better understand in the gas phase the micro-solvation process of biomolecules.

We will present the most recent results on $C_2H_4O_2$ and $C_2H_4O_2-H_2O$.

Concerning glycolaldehyde, the microwave spectrum of one conformer was measured by K.M.MARSTOKK and H.MØLLENDAL *J.Mol.Struc* 16, 259 (1973) and R.A.H.BUTLER et al. *The Astrophysical Journal Supplement Series*, 134,319-321 (2001). The interstellar detection has been made by J.M.HOLLIS et al. *The Astrophysical Journal*, 554:L81-L85 (2001).

Meanwhile four stable conformations have been optimized for glycolaldehyde by *ab initio* calculations by T. Ratajczyk et al. *J. Chem. Phys.A* 108, 2758(2004).

Consequently we have performed additional calculations to obtain rotational constants and electric dipole moments. Our spectrum, recorded in the region 6-20 GHz, revealed new signals that could be assigned to at least one new conformation.

Concerning the glycolaldehyde-water complex, we have performed *ab initio* calculations at the B3LYP/6-311++G(2df,p) and aug-cc-pVTZ levels of the theory. Six stable conformers have been found.

Relative energies have been calculated at the G3MP2B3 level, the most stable structure is stabilized by two inter-molecular hydrogen bonds. The geometry optimization as well as the analysis of the experimental spectrum (6-20GHz) are in progress.

This work is supported by grants IDRIS number 041715 and 051715.