EXPERIMENTAL STUDY OF THE Cs_2 TRIPLET STATES BY PERTURBATION FACILITATED INFRARED-INFRARED DOUBLE RESONANCE AND TWO-PHOTON EXCITATION SPECTROSCOPY

FENG XIE, DAN LI, LI LI, *Department of Physics and Key Laboratory of Atomic and Molecular Nanosciences, Tsinghua University,Beijing 100084, China*; VLADIMIR B. SOVKOV, VALERY S. IVANOV, *V. A. Fock Institute of Physics, St. Petersburg State University, 1 Ulyanovskaya Street, Petrodvorets, St. Petersburg 198504, Russia*; and SYLVIE MAGNIER, *Laboratoire de Physique des Atomes, Lasers, Molećule et Surfaces (PALMS), CNRS et Universite´ RENNES I (UMR 6627), Campus de Beaulieu, Bt 11B 35042 Cedex, France*.

Fifty $3^3\Sigma_g^+$ $v = 2-9 \leftarrow X^1\Sigma_g^+$ two-photon transitions of Cs_2 have been observed in the infrared wavelength. Vibrational resolved fluorescence into both bound levels and the continuum of the $a^3\Sigma_u^+$ state has been recorded with monochromator. Calculated $3^3\Sigma_g^+ \rightarrow a^3\Sigma_u^+$ fluorescence spectra are compared with observed spectra.
With two infrared diode lasers,totally 43 ro-vibrational levels of the Cs_2 $3^3\Pi_g$ state have been observed by perturbation facilitated infrared-infrared double resonance spectroscopy. Rotationally resolved fluorescence to the $a^3\Sigma_u^+$ state was also recorded.