SYNCHROTRON FAR INFRARED SPECTROSCOPY: HIGHER RESOLUTION AND LONGER WAVELENGTHS AT THE CANADIAN LIGHT SOURCE

A.R.W. McKELLAR, *Steacie Institute for Molecular Sciences, National Research Council of Canada, Ottawa, ON K1A 0R6, Canada*; D.R.T. APPADOO[a], *Canadian Light Source, 101 Perimeter Road, University of Saskatchewan, Saskatoon, SK S7N 0X4, Canada*.

In order to take full advantage of the high spectral resolution capability of the Bruker IFS125 Fourier transform spectrometer (MOPD = 9.4 m) at the Canadian Light Source far infrared beamline, we have interfaced a large 2 m gas cell which allows longer absorption paths, lower sample pressures, and hence reduced pressure broadening. We are using acrolein (CH_2CHCHO) for testing. Interesting in its own right, this molecule is also of importance in astrophysics, combustion chemistry, and human respiration (cigarette smoke, smog). Our initial CLS work[b] on ν_{12} ($\sim$564 cm^{-1}) and ν_{17} ($\sim$593 cm^{-1}) has now been extended to ν_{18} ($\sim$158 cm^{-1}), $2\nu_{18}$ ($\sim$314 cm^{-1}), ν_{13} ($\sim$323 cm^{-1}), $3\nu_{18}$ ($\sim$469 cm^{-1}), and further vibrational levels.

We experience significant gains in source intensity and signal-to-noise ratio due to the brightness advantage of synchrotron radiation compared to conventional thermal far infrared sources, though source noise (due to mechanical vibrations) remains a concern. Even greater gains at long wavelengths ($<$50 cm^{-1}) may be possible using coherent synchrotron radiation (CSR), achieved by shortening the electron bunch length in the synchrotron storage ring.[c] Initial testing of CSR at CLS has already yielded a recognizable high-resolution spectrum of N_2O pure rotational transitions in the 10 - 20 cm^{-1} range.

---

[a] Current address: Australian Synchrotron, 800 Blackburn Rd, Clayton VIC 3168, Australia.

[b] A.R.W. McKellar, D.W. Tokaryk, L.-H. Xu, D.R.T. Appadoo, and T. May, *J. Mol. Spectrosc.* **241**, 31 (2007).

[c] M. Abo-Bakr, J. Feikes, K. Holldack, P. Kuske, W. B. Peatman, U. Schade, G. Wüstefeld, and H.-W. Hübers, *Phys. Rev. Letters* **90**, 094801 (2003).