

INVESTIGATION OF THE TORSIONAL FAR-INFRARED OVERTONES AND HOT BANDS OF ACETALDEHYDE AND VIBRATIONAL-TORSIONAL-ROTATIONAL INTERACTIONS WITH THE  $\nu_{10}$  FUNDAMENTAL BAND

I. KLEINER, *Laboratoire de Photophysique Moléculaire, Université Paris Sud, 91405 Orsay, France*;  
N. MOAZZEN-AHMADI, *Department of Physics, University of Lethbridge, Lethbridge, Alberta, T1K 3M4, Canada*;  
A. R. W. McKELLAR, *Stracie Institute for Molecular Science, National Research Council, Ottawa, Canada, K1A0R6*;  
J. T. HOUGEN, *Optical Physics Division, National Institute of Standards and Technology, Gaithersburg, MD 20899, USA*.

The torsional overtone spectrum of CH<sub>3</sub>CHO has been recorded with a high resolution (0.003 cm<sup>-1</sup>) Far-Infrared Fourier Transform spectrometer, between 40 and 90 cm<sup>-1</sup> and between 180 and 230 cm<sup>-1</sup>. This spectrum completes the far-infrared spectrum recorded a few years ago from 80 to 180 cm<sup>-1</sup> to study the fundamental torsional band  $v_t = 1-0$  and the two first overtones  $v_t = 2-1$  and  $3-2^a$ . The spectral region between 40 and 90 cm<sup>-1</sup> is very dense and allows us to assign many  $v_t = 2-1$  lines and  $v_t = 3-2$  lines. The region between 180 and 230 cm<sup>-1</sup>, on the other hand, is less congested and allows us to search for weaker lines from the  $v_t = 3-1$  and  $4-2$  overtone bands. Those overtone bands would correspond, in the free-rotor limit, to a change in the free-rotor quantum number  $m$  of  $\pm 3$  and are expected to arise when considering intensity contribution from the second terms in the Fourier expansion of the dipole moment  $\mu_a^{(3)} \cos 3\alpha$  and  $\mu_b^{(3)} \cos 3\alpha$  for the A and E species and also from the first non-zero term of  $\mu_c^{(3)} \sin 3\alpha$  for the E species. The spectrum of the  $\nu_{10}$  band was recorded last year at high resolution with a Fourier Transform spectrometer in the region of 450 to 550 cm<sup>-1</sup> region and partly assigned. A number of strong perturbations arising likely from interactions with the  $v_t = 4$  torsional levels of the ground state were observed. As a first step to investigate doorway states which facilitate passage of energy from the small amplitude manifold to the torsional manifold, the goal of this study is to consider a two-by-two theoretical model involving both  $\nu_{10}$  and the ground vibrational state and to carry out a simultaneous fit of the perturbed lines from  $v_{10}=1$  and from  $v_t=4$  of the vibrational ground state.

---

<sup>a</sup>I. Kleiner, J. Hougen, J-U. Grabow, S. P. Belov, M. Yu. Tretyakov and J. Cosléou