

THE ROTATIONAL SPECTRA AND HYPERFINE CONSTANTS OF ZrS AND ZrO.

SARA A. BEATON AND MICHAEL C. L. GERRY, *Department of Chemistry, The University of British Columbia, 2036 Main Mall, Vancouver, British Columbia, Canada V6T 1Z1.*

The pure rotational spectra of ZrS and ZrO have been recorded using cavity Fourier transform microwave spectroscopy in the frequency range 9-26 GHz. The molecules were generated by laser ablation of a solid Zr rod with 0.05% of H₂S or O₂, respectively, in either argon or neon.

For ZrS, this is the first measurement of the pure rotational spectrum of the X¹Σ⁺ state. Rotational transitions in the ground vibrational state were observed for all five Zr³²S isotopomers and for the ⁹⁰Zr³⁴S isotopomer in natural abundance. Transitions in the first excited vibrational state were measured for the 3 most abundant isotopomers and a value for the equilibrium bond length calculated. For the ⁹¹Zr³²S isotopomer nuclear hyperfine structure due to the zirconium nucleus was observed and values for $eQq(^{91}\text{Zr})$ and $C_I(^{91}\text{Zr})$ were determined.

The microwave spectra of the three most abundant isotopomers of Zr¹⁶O had been observed previously (1). The new results from the present work are the observations of the $J = 1 - 0$ transitions for the remaining two Zr¹⁶O isotopomers and three Zr¹⁸O isotopomers. In addition, the $v = 1 \quad J = 1 - 0$ transition was measured for ⁹⁰Zr¹⁶O, allowing an accurate determination of r_e . ⁹¹Zr nuclear hyperfine structure was observed for ⁹¹Zr¹⁶O and values of the hyperfine constants were determined. The hyperfine parameters for the two molecules are interpreted in terms of the electronic structure and chemical bonding.

(1) R. D. Suenram, F. J. Lovas, G. T. Fraser and K. Matsumura, *J. Chem. Phys.* **92**, 4725-4733 (1990).