HIGH-RESOLUTION LASER SPECTROSCOPY OF $A^3\Pi_{1u} \leftarrow X^1\Sigma_g^+$ SYSTEM OF I_2

N. NISHIMIYA, T. YUKIYA, O. OHSHIMA, AND M. SUZUKI, *Dept. of Electronic Engineering, Tokyo Institute of Polytechnics, Iiyama 1583, Atsugi City, 243-0297 Kanagawa, Japan*.

The molecular constants for the A-state of I_2 calculated by the NDE method have been reported by D.R.T.Appadoo *et al.*[1] Yukiya *et al.* measured the Q-branch lines up to $v' = 45$ using a titanium sapphire ring laser, where the hyperfine components were partially resolved at the higher vibrational bands than those of $v' = 35$, and determined the spectroscopic constants.[2] In this work we report the line splittings of the P and R-branches caused by the nuclear quadrupole coupling effect. The 23 bands of $(v' = 15 - 19) \leftarrow (v'' = 3)$, $(v' = 18 - 30) \leftarrow (v'' = 4)$, and $(v' = 25 - 29) \leftarrow (v'' = 5)$ were newly assigned. The unperturbed line positions were obtained using the high J approximation.
The spectroscopic constants of T_v, B_{vf}, D_v, H_v, and q_v of the A-state determined by the aid of the Dunham parameters of the X-state[3] will be presented.

1. D.R.T.Appadoo, R.J.LeRoy, P.F.Bernath, S.Gerstenkorn, P.Luc, J.Vergès, J.Sinzelle, J.Chevillard, and Y.D'Aignaux, *J.Chem.Phys.*, **104,** 903 (1996).
2. T.Yukiya, N.Nishimiya, and M.Suzuki, *J.Mol.Spectrosc.*, **182,** 271 (1996).
3. S.Gerstenkorn and P.Luc, *J.Phys.*, **46,** 867 (1985).