

SHORT-WAVELENGTH (30 TO 150 μm) OPTICALLY PUMPED LASER LINES IN A METHANOL LASER

E. C. C. VASCONCELLOS, *Instituto de Física "Gleb Wataghin," Departamento de Eletrônica Quântica, Universidade Estadual de Campinas (UNICAMP), 13083-970 Campinas, SP, Brazil;*

L. R. ZINK and K. M. EVENSON, *Time and Frequency Division, National Institute of Standards and Technology, Boulder, CO 80303-3328.*

A new short-wavelength far-infrared laser^a is used to generate optically pumped far-infrared laser radiation in the wavelength range 30 to 150 μm in methanol. The large-amplitude torsional motion in methanol accounts for its rich far-infrared spectra. Its infrared absorption bands overlap well with the CO₂ laser providing for the generation of strong high-frequency laser lines in the far-infrared. Methanol is an excellent short wavelength laser medium and about 40% of its lines have wavelengths shorter than 150 μm . We have already obtained a four-fold increase in the number of laser lines in the 50 μm region, and are now pushing to increase the density of laser lines with $\lambda < 50 \mu\text{m}$. A list of the strong laser lines observed in methanol, in the range 30 to 150 μm , will be presented in this work.

^aFor detailed description of the laser see paper presented by K.M. Evenson at this meeting.