ROTATIONAL SPECTRA OF THE FREE RADICALS $C_{10}H$, $C_{12}H$, $C_{13}H$, AND $C_{14}H$ IN A SUPERSONIC JET

C. A. GOTTLIEB, M. C. McCARTHY, M. J. TRAVERS, AND P. THADDEUS, *Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138 and Division of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138*.

Four new carbon chain radicals $C_{10}H$, $C_{12}H$, $C_{13}H$, and $C_{14}H$ have been observed in a pulsed supersonic molecular beam with a Fourier transform microwave spectrometer. The radicals were produced in a discharge through a dilute diacetylene/neon mixture in the throat of a supersonic nozzle. All are found to be linear with $^2\Pi$ electronic ground states, and all except $C_{14}H$ have resolved lambda-type doubling. At least 10 rotational transitions, between 6 and 16 GHz, were measured in the lowest spin component — $^2\Pi_{3/2}$ of $C_{10}H$, $C_{12}H$, and $C_{14}H$, and the $^2\Pi_{1/2}$ component of $C_{13}H$. Only three spectroscopic constants in the standard Hamiltonian for a molecule in a $^2\Pi$ state were required to reproduce the spectra to a few parts in $10^7$: an effective rotational constant, a centrifugal distortion constant, and a lambda-type doubling constant. Detection of these highly unsaturated carbon chains establishes that $C_nH$ radicals containing up to 14 carbon atoms are readily produced in a supersonic molecular beam. The relative abundance of $C_nH$ radicals with an *even* number of carbon atoms is fairly constant from $C_6H$ through $C_{12}H$. Although the new radicals are about two orders of magnitude less abundant than $C_4H$, the strong predicted $^2\Pi - ^2\Pi$ electronic transitions may be detectable in a supersonic jet by standard laser spectroscopic techniques.