

THE FORBIDDEN ROTATION AND ROTATION-VIBRATION SPECTRUM OF H_2^+

P. R. BUNKER, *Steacie Institute for Molecular Sciences, National Research Council of Canada, Ottawa, Ontario K1A 0R6, Canada*; R. E. MOSS, *Department of Chemistry, University of Southampton, Southampton SO17 1BJ, U.K.*.

Previous studies of the g/u electronic symmetry breaking interaction between the rotation-vibration levels of the $X^2\Sigma_g^+$ ground electronic state and the $A^2\Sigma_u^+$ first excited electronic state of H_2^+ have focused on explaining the large hyperfine structure observed in some rovibronic transitions between these electronic states^{a, b}. The interaction is an ortho-para interaction between levels of the two electronic states. It gives rise to an interesting new electric dipole spectrum that consists of forbidden rotation and rotation-vibration transitions within the $X^2\Sigma_g^+$ ground state involving levels that are near the dissociation limit^c. Such a spectrum will occur for any homonuclear diatomic ion having nuclei with non-zero spin such as $^3\text{He}_2^+$, Li_2^+ and $^{21}\text{Ne}_2^+$.

^aR. E. Moss, *Chem. Phys. Lett.* **206**, 83 (1993), and references therein.

^bSection 17.7 of the book *Molecular Symmetry and Spectroscopy*, 2nd Edition, by P. R. Bunker and P. Jensen, NRC Research Press, Ottawa 1998. See <http://www.nrc.ca/cisti/journals/41653> for the table of contents and ordering information.

^cP. R. Bunker and R. E. Moss, *Chem. Phys. Lett.* **316**, 266 (2000).