

LASER-INDUCED FLUORESCENCE IN KLi

A. J. ROSS, F. MARTIN, P. CROZET and M. AUBERT-FRÉCON, *Laboratoire de Spectrométrie Ionique et Moléculaire (UMR 5579 CNRS), Bâtiment A. Kastler, Université Lyon I, Domaine Scientifique de la Doua, 69622 Villeurbanne Cedex, France*; P. KOWALCZYK, *Institute of Experimental Physics, Warsaw University, Ul. Hoża 69, 00-681 Warsaw, Poland*; W. JASTRZĘBSKI, *Institute of Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland.*; and A. PASHOV, *Universität Hannover, Institut für Quantenoptik, Welfengarten 1, 30167 Hannover, Germany.*

Laser-induced fluorescence spectra of the $B \rightarrow X$ system of the potassium-lithium molecule have been recorded on a Fourier transform interferometer. The molecules were excited with a tuneable, single-mode cw dye laser operating with Rhodamine 110 or Coumarine 6 dye, and the spectra recorded in the range 12500 - 18500 cm^{-1} at a resolution of 0.05 cm^{-1} . Vibrational levels up to $v = 46$ are observed for the main isotopomer, $^{39}\text{K}^{7}\text{Li}$.

Ground state energies have been fitted to a Dunham polynomial expansion, and also directly to a pointwise potential curve. Both approaches fit the data with a root-mean-square error of 0.005 cm^{-1} . The data define the ground state potential curve accurately out to about 7 Å. The dissociation energy of the ground state D_e is estimated to be $D_e = 6216 \pm 2 \text{ cm}^{-1}$, and the highest vibrational level $v_{max} = 50$ for the main isotopomer.

Very weak transitions around 12600 cm^{-1} are tentatively assigned as bands of the $2^3\Sigma^+ \rightarrow 1^3\Sigma^+$ transition, observed as collisionally induced fluorescence.