

SPECTRAL INHOMOGENEITY AS A DISCRETE, COUNTABLE ENTITY – THE ROLE OF RESIDUAL ORTHO-HYDROGEN MOLECULES ON THE ROVIBRATIONAL SPECTRA OF DOPANTS IN PARAHYDROGEN SOLIDS

MARIO E. FAJARDO and SIMON TAM, *AFRL/PRSP, Bldg. 8451, Propulsion Sciences and Advanced Concepts Division, US Air Force Research Lab, Edwards AFB, CA 93524-7680.*

We report high resolution (0.005 cm^{-1}) infrared (IR) absorption spectra of doped millimeters-thick parahydrogen (pH_2) solids with intentionally elevated residual orthohydrogen (oH_2) concentrations. In addition to the weakly-hindered rotor spectrum of isolated H_2O molecules, $\text{H}_2\text{O}/\text{oH}_2/\text{pH}_2$ samples exhibit a series of sharp (0.03 cm^{-1} FWHM) absorption peaks in the 1593 to 1597 cm^{-1} region which we show are due to $\text{H}_2\text{O}-\text{oH}_2$ clusters. We have also identified CO_2-oH_2 clusters as the source of the very sharp (0.007 cm^{-1} FWHM) fine structure in IR spectra of $\text{CO}_2/\text{oH}_2/\text{pH}_2$ solids.^a The gradual evolution of the CO_2-oH_2 features with increasing oH_2 concentration raises the interesting notion of spectral inhomogeneity as a discrete, countable entity.

^aS. Tam and M.E. Fajardo, *Low Temp. Phys.* v26, p653 (1998).