

HIGH RESOLUTION LASER SPECTROSCOPY OF THE TITANIUM MONOHALIDES TiCl AND TiBr.

A. G. ADAM and SCOTT HOPKINS, *Department of Chemistry, University of New Brunswick, Fredericton, NB, Canada E3B 6E2.*

High resolution laser spectra of TiCl and TiBr have been taken in the blue region of the spectrum. The titanium monohalide molecules were produced by laser vaporization of a Ti rod followed by reaction with either CHCl_3 or CH_3Br in a pulsed supersonic jet source. The TiCl molecule was recently studied by FT spectroscopy^a where the electronic transition was identified as ${}^4\Gamma - {}^4\Phi$ consistent with that of TiF. In the paper, band-by-band case (c) rotational analyses of the (0,0) ${}^4\Gamma_{5/2} - {}^4\Phi_{3/2}$, ${}^4\Gamma_{7/2} - {}^4\Phi_{5/2}$ and ${}^4\Gamma_{9/2} - {}^4\Phi_{7/2}$ subbands were given. The authors observed no subbands which would provide the information crucial to determining the spin-orbit spacing. We have observed this same TiCl electronic transition with our apparatus and report here the observation of the (0,0) ${}^4\Gamma_{5/2} - {}^4\Phi_{5/2}$ and ${}^4\Gamma_{7/2} - {}^4\Phi_{7/2}$ subbands, in addition to those given above, and provide a case (a) molecular analysis. The same electronic transition for TiBr has also been recorded. Although the spectrum is much more congested than that of TiCl, we have managed to assign a number of the subbands and are also able to provide a case (a) analysis for this molecule.

^aT. Imajo, D. Wang, K. Tanaka, and T. Tanaka, *J. Mol. Spectrosc.* **203**, 216 (2000).