

## BROADENING AND SHIFT COEFFICIENTS IN THE $^{12}\text{C}^{16}\text{O}_2$ AND $^{13}\text{C}^{16}\text{O}_2$ LASER BANDS

D. CHRIS BENNER, V. MALATHY DEVI, *Department of Physics, The College of William and Mary, Box 8795, Williamsburg, VA 23187-8795*; M. A. H. SMITH, C. P. RINSLAND, *Atmospheric Sciences, NASA Langley Research Center, Mail Stop 401A, Hampton, VA 23681-2199*; L. R. BROWN, *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109*; and M. DULICK, *National Solar Observatory, P. O. Box 26732, Tucson, AZ 85726-6732*.

In a previous study<sup>a</sup> we had reported air- and N<sub>2</sub>-broadening and pressure-induced shift coefficients for 46 rovibrational transitions in the  $^{12}\text{C}^{16}\text{O}_2$   $00^01-(10^00-02^01)_I$  and  $00^01-(10^00-02^01)_{II}$  laser bands near 10  $\mu\text{m}$ . These parameters were determined from spectra recorded with the McMath-Pierce Fourier transform spectrometer (FTS) of the National Solar Observatory on Kitt Peak, Arizona. We now report similar measurements of air- and N<sub>2</sub>-broadening and shifts for the two  $^{13}\text{C}^{16}\text{O}_2$  laser bands, plus new determinations of self-broadening and shifts in the  $^{12}\text{C}^{16}\text{O}_2$  laser bands. These results were obtained from simultaneous analysis of 30 long-path absorption spectra using a multispectrum nonlinear least-squares technique<sup>b</sup>. In addition to the 10 spectra of air- and N<sub>2</sub>-broadened natural CO<sub>2</sub> analyzed previously, we have included 20 new spectra recorded at room temperature and 0.005  $\text{cm}^{-1}$  resolution using the McMath-Pierce FTS. These additional spectra include two low pressure, four air-broadened and four N<sub>2</sub>-broadened spectra obtained with a 90% <sup>13</sup>C-enriched CO<sub>2</sub> sample and 10 self-broadened spectra obtained with a high purity natural CO<sub>2</sub> sample. The self-broadened spectra were recorded with pathlengths of 73, 97, 193 and 433 m and CO<sub>2</sub> pressures varying between 3 Torr and 400 Torr. The  $^{13}\text{CO}_2$  spectra were obtained with a pathlength of 84 m with sample pressures  $\geq 4$  Torr for the low-pressure spectra and from 100 to 500 Torr for the air- and N<sub>2</sub>-broadened mixtures. By combining the spectra of  $^{12}\text{CO}_2$  and  $^{13}\text{CO}_2$  in the same fit we were able to obtain a consistent set of line parameters for both molecules. The results obtained for the  $^{12}\text{CO}_2$  and  $^{13}\text{CO}_2$  laser bands will be compared with each other, with the values in the HITRAN database<sup>c</sup>, and with available values reported in the literature.

<sup>a</sup>V. Malathy Devi, D. Chris Benner, M. A. H. Smith and C. P. Rinsland, *JQSRT* **59**, 137-149 (1998).

<sup>b</sup>D. Chris Benner, C. P. Rinsland, V. Malathy Devi, M. A. H. Smith and D. Atkins, *JQSRT* **53**, 705-721 (1995).

<sup>c</sup>L. S. Rothman et al., *JQSRT* **60**, 665-710 (1998).