

## TWO-ROTOR CALCULATIONS FOR MOLECULES WITH $C_2$ OR $C_s$ SYMMETRY

PETER GRONER, *Department of Chemistry, University of Missouri - Kansas City, Kansas City, MO 64110.*

An effective rotation-internal motion Hamiltonian for molecules with two periodic internal motions <sup>a</sup> has been applied recently to dimethyl ether <sup>b</sup> and acetone <sup>c</sup> whose equilibrium structures have  $C_{2v}$  symmetry. The program used to analyze their rotational spectra has been modified to allow treatment of two-rotor molecules with lower symmetry (equivalent rotor problems for molecules with  $C_s$  and  $C_2$  symmetry and nonequivalent rotor problems with and without a plane of symmetry). It has been used to analyze the microwave literature data of  $\text{CH}_3\text{SSCH}_3$  ( $C_2$  symmetry, ground state) <sup>d</sup>,  $\text{CH}_3\text{CHFCH}_3$  ( $C_s$  symmetry, ground and two excited states) <sup>e</sup>, and  $\text{CH}_3\text{OSiH}_3$  (nonequivalent,  $C_s$  symmetry, ground state) <sup>f</sup>. The experimental data have been fitted to experimental precision by the nonlinear least-squares method to determine spectroscopic parameters (rotational constants, centrifugal distortion constants and tunneling parameters). Applications to other molecules are in progress.

---

<sup>a</sup>P. Groner, J. Chem. Phys. 107, 4483 (1997)

<sup>b</sup>P. Groner, S. Albert, E. Herbst and F. C. De Lucia, Astrophys. J. 500, 1059 (1998)

<sup>c</sup>P. Groner, S. Albert, E. Herbst, F. C. De Lucia, F. J. Lovas, J. C. Pearson and B. J. Drouin, Astrophys. J. (submitted)

<sup>d</sup>M. Meyer, J. Mol. Struct. 273, 99 (1992)

<sup>e</sup>M. Meyer and H. Dreizler, Z. Naturforsch. A 43, 138 (1988); J. Mol. Spectrosc. 148, 310 (1991)

<sup>f</sup>C. D. LeCroix, R. F. Curl, P. M. McKinney and R. J. Myers, J. Mol. Spectrosc. 53, 250 (1972)