

TORSIONAL ANALYSIS OF 2-BUTYNOL

RANGANATHAN SUBRAMANIAN, *Department of Chemistry, Wesleyan University, Middletown, CT 06459*; ROBERT K. BOHN, *Department of Chemistry, University of Connecticut, Storrs, CT 06269*; and STEWART E. NOVICK, *Department of Chemistry, Wesleyan University, Middletown, CT 06459*.

Microwave spectroscopic studies of substituted propynes, $\text{CH}_3\text{-C}\equiv\text{C-X}$, have increased our understanding of the nature and origin of the potential barriers for internal rotation. Specifically, the potential barriers have been measured for the groups, $-\text{CD}_3^a$, $-\text{CH}_2\text{Cl}^b$, $-\text{SiH}_3^c$, and $-\text{COF}^d$, bonded to the propynyl fragment, with reported 3-fold barriers of 5.62(16), 10.05(9), 3.77(70) and 2.20(12) cm^{-1} respectively. We report here a study of the rotational spectrum of 2-butyne-1-ol, $\text{CH}_3\text{-C}\equiv\text{C-CH}_2\text{OH}$. 2-butyne-1-ol has two low energy rotational conformers with the OH group either anti or gauche with respect to the propynyl moiety. The study of 2-butyne-1-ol was performed with a pulsed-jet, Fourier Transform Microwave Spectrometer. The rotational spectrum of one species including tunneling splittings has been assigned and is consistent with the gauche conformer. A preliminary value of the 3-fold barrier is 12.5(4) cm^{-1} . Unassigned transitions in the spectrum may be due to the other conformer.

^aJ. Nakagawa, M. Hayashi, Y. Endo, S. Saito, and E. Hirota *J. Chem. Phys.* **80**, 5922 (1966)

^bV.M. Stolwijk and B.P. van Eijck *J. Mol. Spectrosc.* **124**, 92 (1987)

^cJ. Nakagawa, K. Yamada, M. Bester, and G. Winnewisser *J. Mol. Spectrosc.* **110**, 74 (1985)

^dK. D. Hensel and M. C. L. Gerry *J. Chem. Soc. Faraday Trans.* **90**, 3023 (1992)