

FOURIER TRANSFORM MICROWAVE SPECTROSCOPY OF N-METHYLPROPIONAMIDE

YOSHIYUKI KAWASHIMA, *Department of Applied Chemistry, Kanagawa Institute of Technology, Atsugi, Kanagawa 243-0292, JAPAN*; RICHARD D. SUENRAM, *National Institute of Standards and Technology, Gaithersburg, MD 20899*; and EIZI HIROTA, *The Graduate University for Advanced Studies, Hayama, Kanagawa 240-0193, JAPAN*.

In order to clarify the dynamical behavior of the peptide bond, we have undertaken a systematic study of 'peptide molecules', which consist of (a) peptide bond(s) with internal-rotation groups at the both ends of the bond(s). In the present investigation we focused attention to the molecule (NMPA) shown in the title, which has an ethyl group (CH_3 of which is referred to as C- CH_3) at the carbonyl side and a methyl group (called N- CH_3) at the amide side, and aimed at unveiling how the two CH_3 groups interact with each other through the peptide bond. We have derived a rotational Hamiltonian including the two CH_3 internal rotations, and have treated the C- CH_3 internal rotation by a conventional PAM, while applying a more sophisticated approach to the N- CH_3 internal rotation. NMPA may be regarded to belong to group G_{18} , even if its skeleton executes large-amplitude 'out-of-plane' motions. The group consists of 6 species: A_1, A_2, E_1, E_2, E_3 , and E_4 . We have observed and analyzed A_1 (or A_2) and E_2 spectra, but have not detected any lines due to the first excited state of the $\text{CH}_3\text{CH}_2\text{-CO}$ torsion of A_2 symmetry, indicating that the internal-rotation splitting is quite large. The potential barrier to the C- CH_3 internal rotation was determined to be 799 cm^{-1} , which may be compared with that of N- CH_3 of about 81 cm^{-1} . The coupling between the two CH_3 's is being analyzed by observing E_1, E_3 , and E_4 spectra.