

INFRARED PHOTODISSOCIATION SPECTROSCOPY OF MASS-SELECTED METAL ION-BENZENE COMPLEXES

M. A. DUNCAN, DEPARTMENT OF CHEMISTRY, UNIVERSITY OF GEORGIA, ATHENS, GA. 30602; D. van HEIJNSBERGEN, G. von HELDEN, and G. MEIJER, F. O. M. LABORATORY FOR PLASMAPHYSICS RIJNHUISEN, EDINSONBAAN 14, NL-3439 MN NIEUWEGEIN, THE NETHERLANDS.

Metal ion-benzene complexes of the form $M^+(\text{benzene})_{1,2}$ ($M=\text{V, Ti, Al}$) are produced by laser vaporization in a pulsed nozzle source, mass-selected in an ion-trap mass spectrometer and excited in the infrared with a free electron laser. Multiphoton dissociation occurs by the elimination of benzene molecules. The photofragment yield versus wavelength produces IR resonance-enhanced multiphoton photodissociation (IR-REMPD) spectroscopy for these complexes. Vibrational bands in the $600\text{-}1800\text{cm}^{-1}$ region are characteristic of the benzene molecular moiety perturbed by the metal bonding. Experimental data are compared to the IR spectra derived from density functional calculations. Vibrational patterns in $V^+(\text{C}_6\text{H}_6)$ indicate that the metal is bound in an η^6 π -bonding configuration, while $V^+(\text{C}_6\text{H}_6)_2$ is a sandwich. Trapped-ion IR-REMPD is a general method to access the vibrational spectra of novel organometallic complexes.