

THE PERMANENT ELECTRIC DIPOLE MOMENTS FOR THE  $A^2\Pi$  AND  $B^2\Sigma^+$  STATES AND THE HYPERFINE INTERACTIONS IN THE  $A^2\Pi$  STATE OF LANTHANUM MONOXIDE

WILTON L. VIRGO and TIMOTHY C. STEIMLE, *Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ, 85287-1604.*

The optical Stark effect in the (0,0)  $A^2\Pi_{3/2} - X^2\Sigma^+$ ,  $A^2\Pi_{1/2} - X^2\Sigma^+$  and  $B^2\Sigma^+ - X^2\Sigma^+$  band systems of a lanthanum monoxide, LaO, supersonic molecular beam sample have been analyzed to produce permanent electric dipole moments,  $\mu$ , for the  $A^2\Pi_{3/2}$ ,  $A^2\Pi_{1/2}$  and  $B^2\Sigma^+$  states of 1.89(6)D, 2.44(2)D and 0.2(1)D, respectively. The ground state dipole moment was determined some time ago<sup>1</sup> to be 3.207(11)D. Fine structure splitting in the field free  $A^2\Pi_{3/2} - X^2\Sigma^+$  and  $A^2\Pi_{1/2} - X^2\Sigma^+$  spectra was analyzed to produce the magnetic hyperfine spectroscopic parameters  $a = 233(4)$  MHz,  $c = -261(12)$  MHz, and  $d = 410(4)$  MHz. The Group IIIA monoxides, ScO, YO and LaO are prototypical examples of transition metal-oxide bonded molecules. This work on LaO has resulted in the first complete set of dipole moment measurements for the  $X^2\Sigma^+$ ,  $A^2\Pi_{1/2}$ ,  $A^2\Pi_{3/2}$  and  $B^2\Sigma^+$  states for any of these monoxides. The permanent electric dipole moments for the low-lying states of LaO are exceedingly small, given that they are conventionally viewed as having a  $La^{+2}O^{-2}$  charge distribution and exhibiting a large variation amongst the four states.

1. R.F. Seunram, F.J. Lovas, G.T. Fraser and K. Matsumura, J. Chem. Phys. 92, 4724 (1990).