

INVESTIGATION OF BORN-OPPENHEIMER BREAKDOWN IN HCl/DCI BY THRESHOLD ION-PAIR PRODUCTION SPECTROSCOPY (TIPPS)

TODD C. MELVILLE, QICHI J. HU, JOHN W. HEPBURN, *Department of Chemistry, University of British Columbia, Vancouver, BC, Canada V6T 1Z1.*

The spectroscopic technique Threshold Ion-Pair Production Spectroscopy (TIPPS)^a is applied for the first time to the isotopomers HCl/DCI. From the high-resolution TIPP spectra, the difference between the equilibrium bond dissociation energies of the isotopomers can be measured to better than 1 cm^{-1} , and therefore an experimental estimate of the Born-Oppenheimer breakdown in the long range limit of the ground state potential curve is obtained. The experimental result from the present work is compared to a recent determination of the Born-Oppenheimer breakdown in HCl by Coxon and Hajigeorgiou.^b If time permits, similar work that is in progress on HF/DF will also be discussed.

^aJ. D. D. Martin, J. W. Hepburn, *Phys. Rev. Lett.* 79, 3154 (1997).

^bJ. A. Coxon, P. G. Hajigeorgiou, *J. Mol. Spectrosc.* 203, 49 (2000).