

ROTATIONAL SPECTRA AND *AB INITIO* POTENTIAL ENERGY SURFACE OF THE H<sub>2</sub>-OCS VAN DER WAALS COMPLEX

KELLY HIGGINS, Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455;  
ZHENHONG YU, and WILLIAM KLEMPERER, Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138.

The four-dimensional intermolecular potential energy surface for the H<sub>2</sub>-OCS complex was obtained at the MP4 level. The potential gives a T-shaped global minimum with a distance of 3.2 Å between H<sub>2</sub> and OCS, in which H<sub>2</sub> is nearly parallel to OCS. Bound state calculations of *para*H<sub>2</sub>-OCS give predicted rotational constants of A = 22652, B = 5994, and C = 4611 MHz, in good agreement with the measured results from high-resolution infrared studies.<sup>a</sup> The calculated binding energy of *para*H<sub>2</sub>-OCS is 75 cm<sup>-1</sup>, almost four times greater than that of He-OCS.<sup>b</sup> Preliminary bound state calculations of *ortho*H<sub>2</sub>-OCS predict binding energies of 99 cm<sup>-1</sup> for the Σ state and 64 cm<sup>-1</sup> for the Π state.

*a*-type rotational transitions of two isotopomers of the *ortho*H<sub>2</sub>-OCS complex were observed between 9 to 31 GHz. The spectral constants of *ortho*H<sub>2</sub>-OC<sup>32</sup>S are (B+C)/2 = 5113.372(23), (B-C)/2 = 580.337(40), and D<sub>J</sub> = 2.118(2) MHz. The observed 1<sub>01</sub>-0<sub>00</sub> transition is very close to the expectation of *para*H<sub>2</sub>-OCS. This shows that the ground state of the complex is a Σ bound state with the T-shaped geometry. The H<sub>2</sub> nuclear spin dipole-dipole coupling constant d<sub>H H</sub><sup>c</sup> is 14.4(1) kHz which indicates a zero-point energy averaged angle of 45° of H<sub>2</sub> with respect to the molecular *a* axis. Preliminary spectra of D<sub>2</sub>-OCS show the existence of both *ortho*D<sub>2</sub>-OCS and *para*D<sub>2</sub>-OCS complexes.

---

<sup>a</sup>Jian Tang and A. R. W. McKellar, J. Chem. Phys. 116, 646 (2002).

<sup>b</sup>Kelly Higgins and William Klemperer, J. Chem. Phys. 103, 1138 (1999).

<sup>c</sup>N. F. Ramsey, Molecular beam (Oxford University, London, 1953).