

FOURIER TRANSFORM AND INTRACAVITY LASER SPECTROSCOPY OF NICKEL CHLORIDE: IDENTIFICATION OF THE X $^2\Pi_{1/2}$ STATE

LEAH OBRIEN, COREY RICE, BARTHEMEAUS OWEN, and TENEIL KELLERMAN, *Department of Chemistry, Southern Illinois University, Edwardsville, IL 62026-1652*; JAMES OBRIEN and HONG CAO, *Department of Chemistry, University of Missouri, St Louis, MO 63121-4499*.

The near infrared electronic transition of NiCl occurring in the region of 11900 cm^{-1} , also known as System I, has been recorded with rotational resolution by Fourier transform emissions spectroscopy and intracavity laser absorption spectroscopy. This band has been identified as the $[12.3] \ ^2\Sigma^+ - X \ ^2\Pi_{1/2}$ transition. The molecular constants for the newly identified X $^2\Pi_{1/2}$ state are (in cm^{-1}): $T_0=385.666(2)$, $B_0=0.180778(2)$, $D_0=1.652(3)\times 10^{-7}$, $p=0.8391(1)$, $p_D=3.90(7)\times 10^{-6}$. Results of the analysis will be discussed. The identification of the low-lying X $^2\Pi_{1/2}$ state completes the analyses of low-lying electronic states of NiCl that correlate to the $\text{Ni}^+ 3d^9$ electron configuration. Progress on using a 5×5 supermultiplet to describe the low-lying states of NiCl will be reported.