

HIGH-RESOLUTION SPECTROSCOPY OF THE (0,2) AND (6,9) BANDS OF THE $B^4\Sigma_u^- \leftarrow X^4\Sigma_g^+$ TRANSITION OF C_2^+

C. G. TARSITANO, C. F. NEESE, and T. OKA, *Department of Chemistry, Department of Astronomy and Astrophysics, and the Enrico Fermi Institute, University of Chicago, Chicago, IL 60637.*

The first electronic spectrum of the molecular ion C_2^+ was recorded in 1984 by O'Keefe *et. al.* in a beam of mass selected ions^a. Three years later, Maier and coworkers did an extensive analysis on several bands of the $B \leftarrow X$ system^{bcd}. While their spectra showed several perturbations, the low resolution of their experiment hindered the analysis of the perturbation and the quartet splitting. In 1993, Zackrisson and Royen obtained a high resolution spectrum of the (0,1) band and determined the molecular constants and perturbation parameters^e.

In the work that follows, velocity modulation with heterodyne detection has been used to record the (0,2) and (6,9) bands of the $B \leftarrow X$ system of C_2^+ . The (0,2) band has been rotationally analyzed and the spectrum shows perturbations which are attributed to an interaction between the $B^4\Sigma_u^-$ and the $2^2\Pi_u$ states. Molecular constants and perturbation parameters have been obtained.

^aA. O'Keefe, R. Derai, and M. T. Bowers, *Chem. Phys.* **91**, 161 (1984)

^bM. Rösslein, M. Wyttenbach, and J. P. Maier, *J. Chem. Phys.* **87**, 6770 (1987)

^cJ. P. Maier and M. Rösslein, *J. Chem. Phys.* **88**, 4614 (1988)

^dF. G. Celil and J. P. Maier, *Chem. Phys. Lett.* **166**, 517 (1990)

^eM. Zackrisson and P. Royen, *J. Mol. Spectrosc.* **161**, 1 (1993)