

## ADDITION OF WATER TO AMMONIA-HYDROGEN HALIDE COMPLEXES: MOVING TOWARDS PROTON TRANSFER

BRIAN E. APPELGATE, JEREMY M. MERRITT, AND ROGER E. MILLER, *University of North Carolina at Chapel Hill, Department of Chemistry, Chapel Hill, NC 27599.*

Proton transfer in the ammonia-hydrogen halides serves as a prototypical acid-base reaction. Understanding the role of water in facilitating these types of reactions is important for both atmospheric chemistry and solution chemistry. Gas-phase and matrix experiments<sup>a</sup> show that the proton remains attached to the halogen in anhydrous complexes for HF, HCl, and HBr. However, recent theoretical calculations<sup>b</sup> have shown that the proton transfer proceeds with the addition of from one to three water molecules, for HBr, HCl, and HF, respectively. To study this reaction we have initiated a set of experiments where we sequentially add water molecules to preformed ammonia-hydrogen halide complexes inside helium droplets. Observation of the vibrational frequencies along with estimates of the dipole moments help to determine if proton transfer has taken place.

---

<sup>a</sup>Barnes, A.; Legon, A., *J. Mol. Struct.*, **1998**, 101.

<sup>b</sup>Snyder, J.; Cazar, R.; Jamka, A.; Tao, F-M, *J. Phys. Chem.*, **1999**, 7719.