

SELECTIVE PHOTO BLEACHING OF CARBON CHAIN ABSORPTIONS IN CRYOGENIC MATRICES

DMITRY STRELNIKOV, ROMAN REUSCH, and WOLFGANG KRÄTSCHEMER, *Max-Planck-Institut für Kernphysik, D-69029 Heidelberg, PO Box 103980, Germany.*

We present results obtained on combined UV-vis photo bleaching and IR spectroscopy performed on the same matrix sample. Using an excimer pumped dye laser we could selectively deplete the species C₉, C₁₁, C₁₃ and C₁₅ in reactive matrices such as solid oxygen by exposures at the appropriate UV-vis peak absorption wavelengths, which usually are close to those in argon matrices^a. The correlated decreases in UV-vis and IR line intensities yield information on the IR active vibrations of such long chains. According to our data linear C₁₅ exhibits major IR absorptions at 1707-1721 cm⁻¹ in oxygen matrices. In the process of photo bleaching we produced as reaction products various higher carbon oxides, most of which seem to be also chain molecules. The characterisation of these species by ¹⁶O-¹⁸O and ¹²C-¹³C isotopic replacement experiments is in progress. We also attempt to assign carbon chains larger than C₁₅.

^aJ.P. Maier; *Chemical Society Reviews*, 1997, 21 (1997)