

INFRARED DIODE LASER SPECTROSCOPY OF THE P=O (ν_2) FUNDAMENTAL BAND OF *CIS*-HOPO

PAUL J. O'SULLIVAN, PETER A. HAMILTON and PAUL B. DAVIES, *Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, U.K.*

The P=O (ν_2) stretching fundamental of the transient molecule *cis*-HOPO has been measured using high resolution infrared diode laser absorption spectroscopy. The molecule was formed in a flow system by reacting white phosphorus vapour and the products of a microwave discharge through a mixture of hydrogen and oxygen. Over 300 lines have been assigned between 1242cm^{-1} and 1267cm^{-1} . The band has the structure of an a-type transition of an asymmetric rotor. Most of the assigned lines were included in a least squares fit using Watson's A-reduced Hamiltonian. The band origin is $1258.540021(12)\text{cm}^{-1}$. The rotational constants are in satisfactory agreement with those from Density Functional Theory and with those predicted from a comparison with the nitrogen analogue, *cis*-HONO:

$$A_0 = 1.255036(55), B_0 = 0.3188211(99), C_0 = 0.2533085(91)$$

$$A_1 = 1.250424(55), B_1 = 0.3181419(96), C_1 = 0.2526089(96)$$

Many of the remaining, unfitted lines are perturbed, particularly around $K_a=7$. The source of the perturbation will be considered.