

## NEAR INFRARED LASER SPECTROSCOPY OF NiI

W. S. TAM, JIANJUN YE and A. S-C. CHEUNG, *Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong.*

Laser induced fluorescence spectrum of NiI in the near infrared region 714 - 770 nm has been recorded and analyzed. Eleven bands belonging to four electronic transition systems have been observed: the (0,0), (1,0), and (2,0) bands of [13.3]  $^2\Sigma^+$  - A  $^2\Pi_{3/2}$  system; the (0,0), (1,1) and (0,1) bands of [13.9]  $^2\Pi_{3/2}$  - X  $^2\Delta_{5/2}$  system; the (0,0) and (1,0) bands of [13.9]  $^2\Pi_{3/2}$  - A  $^2\Pi_{3/2}$  system; and the (0,0), (1,1) and (0,1) bands of the [14.0]  $^2\Phi_{7/2}$  - X  $^2\Delta_{5/2}$  system. Spectra of isotopic molecules confirmed the vibrational quantum number assignment of the observed bands. Least squares fit of observed rotational lines yielded accurate molecular constants for the [13.3]  $^2\Sigma^+$ , [14.0]  $^2\Phi_{7/2}$  and the A  $^2\Pi_{3/2}$  electronic states of NiI. Hot band transition from the  $v = 1$  level of the X  $^2\Delta_{5/2}$  state gives  $\Delta G_{1/2} = 276.67 \text{ cm}^{-1}$ . The A  $^2\Pi_{3/2}$  state was determined to be only  $163.85 \text{ cm}^{-1}$  above the X  $^2\Delta_{5/2}$  state.