

SPECTRAL ANALYSIS OF THE BENDING LEVELS OF ACETYLENE CATION

YUNG-CHING CHOU and YEN-CHU HSU, *INSTITUTE OF ATOMIC AND MOLECULAR SCIENCES, ACADEMIA SINICA, P.O. BOX 23-166, TAIPEI 107, TAIWAN, R. O. C.*

Twenty-seven vibronic levels ($v_4=0-6$, $v_5=0-2$, $K=0-3$) of the ground electronic state of acetylene cation have been previously studied by 1+1' two-color pulsed-electric field induced zero-kinetic energy (PFI-ZEKE) photoelectron spectroscopy^a. These levels, rotationally resolved, are about 0-3520 cm^{-1} above the zero-point energy level. Previously, only the Renner-Teller effects of the trans- and cis-bending vibration and the vibrational anharmonicity were included in the effective Hamiltonian^b; one standard deviation of the fit was 4.1 cm^{-1} , about six times of the measurement uncertainties. Large discrepancies between the observed and calculated frequencies were identified at combination levels. In this fit, effects such as vibrational ℓ -type doubling and second-order anharmonic interaction were also considered due to the near degeneracy of the trans- and cis-bending vibration. The results of the fit are satisfactory; most of the vibronic bands can be fitted within one standard deviation of 1.0 cm^{-1} , except the lower component of ($v_4=0$, $v_5=1$, $K=0$). The results of the fit will be presented and few observed vibronic levels, yet unassigned, will be discussed.

^aShuenn-Jiun Tang, *Ph. D. thesis*, Chemistry Department, National Taiwan University, Taiwan, R. O. C., 1999.

^bJ. M. Brown, *J. Mol. Spectrosc.* **68**,412(1977).