

NEW ANALYSIS OF THE ν_1 , ν_5 , $2\nu_4$, $\nu_4+\nu_6$, $2\nu_6$, $\nu_3+\nu_4$, $\nu_3+\nu_6$, $\nu_2+\nu_4$, $2\nu_3$, $\nu_2+\nu_6$, AND $\nu_2+\nu_3$ BANDS OF FORMALDEHYDE $\text{H}_2^{12}\text{C}^{16}\text{O}$. LINE POSITIONS AND INTENSITIES IN THE 3.6 μm SPECTRAL REGION.

A. PERRIN, *Laboratoire Inter Universitaire des Systemes Atmosphériques, CNRS, Université Paris 12, 61 Av du General de Gaulle, 94010 Créteil Cedex France*; A. VALENTIN, *Laboratoire de Physique Moléculaire et Applications, C.N.R.S., Université Pierre et Marie Curie, Tour 13, 4 Place Jussieu, 75252 Paris, Cedex 05, France*; L. DAUMONT, *Groupe de Spectroscopie Moléculaire, Faculté des Sciences, BP 347, 51062 Reims, Cedex, France*.

This work is mainly motivated by the tropospheric importance of formaldehyde. The 3.6 μm region is indeed widely used for remote sensings of this molecule in the troposphere by various ground based or portable techniques (diode laser, difference frequency generation (DFG), Fourier transform, and by mid-infrared cavity leak-out spectroscopies). On the other hand the line list which is presently available in HITRAN ^a for this molecule in this spectral range is of poor quality. Using Fourier transform spectra recorded at high resolution at LPMA Paris and at GSMA Reims, it has been possible to perform a new study of the formaldehyde absorption spectra in the 2400-3400 cm^{-1} spectral range. This analysis was started using the results obtained by previous studies performed in the same spectral regions ^{b c}. The analysis of the strongest bands, namely ν_1 and ν_5 (symmetric and antisymmetric stretching modes) was complicated by the existence of numerous dark overtones or combination bands namely $2\nu_4$, $\nu_4+\nu_6$, $2\nu_6$, $\nu_3+\nu_4$, $\nu_3+\nu_6$, $\nu_2+\nu_4$, $2\nu_3$, $\nu_2+\nu_6$, and $\nu_2+\nu_3$. The Hamiltonian model accounts for the various A-type, B-type and C-type Coriolis-type resonances and of the Fermi or anharmonic resonances which perturb the energy levels of the 4^2 , 4^1+6^1 , 6^2 , 3^1+4^1 , 3^1+6^1 , 1^1 , 5^1 , 2^1+4^1 , 3^2 , 2^1+6^1 , and 2^1+3^1 vibrational states. This is also the case for the line intensity calculations which allow one to reproduce satisfactorily the line by line intensity measurements available in the literature.

^aL. H. Rothman, et al. <http://cfa-www.harvard.edu/HITRAN/>

^bL. R. Brown, R. H. Hunt, and A. Pine, *J. Mol. Spectrosc.* **75**, 406 (1979).

^cF. Ito, T. Nakanaga, and H. Takeo, *Spectrochimica Acta.* **50A**, 1397 (1994).