


MICROWAVE SPECTROSCOPY MEASUREMENTS OF THE GAS PHASE STRUCTURE OF CYCLOPENTADIENYL TUNGSTENTRICARBONYL HYDRIDE ^a

CHANDANA KARUNATILAKA, CHAKREE TANJAROON, KRISTEN S. KECK, AND STEPHEN G. KUKOLICH,
Department of Chemistry, University of Arizona, Tucson, AZ 85721.

Microwave spectra for five unique ¹³C isotopomers of cyclopentadienyltungstentricarbonyl hydride were measured using a Flygare-Balle type microwave spectrometer system. The new rotational constants were combined with the previously obtained rotational constants for normal and deuterium analogues for various tungsten isotopes. A total of fifty-seven rotational constants were acquired and used in the least squares fit to obtain the gas phase structure of this d⁴ - metal mono hydride complex. The results from the structural fit yielded the W-H bond length: $r_o(\text{W-H}) = 1.788(20) \text{ \AA}$, which agrees very well with the previously reported Kraitchman value of $r_s(\text{W-H}) = 1.79(4) \text{ \AA}$. The present study also yielded the distance from tungsten to the centroid distance of the C₅H₅ ring: $r(\text{W-Cp}) = 2.02(1) \text{ \AA}$, the ring radius of Cp: $r(\text{Cp}) = 1.2016(4) \text{ \AA}$, the average bond length from tungsten to the carbonyl carbon, $r(\text{W-CO}) = 1.981(14) \text{ \AA}$, and the average cyclopentadienyl C-C bond length of $1.421(4) \text{ \AA}$. Results obtained from the structural fit are in much closer agreement with the Kraitchman values than with the DFT results.



^aSupported by THE NATIONAL SCIENCE FOUNDATION - CHE0304969