

## LINE INTENSITIES MEASUREMENTS IN $^{13}\text{C}^{16}\text{O}_2$ AND THEIR TREATMENT USING THE EFFECTIVE OPERATORS APPROACH

V. I. PEREVALOV, S. A. TASHKUN, *Institute of Atmospheric Optics, Russian Academy of Sciences, Siberian Branch, 1 Akademicheskii Avenue, 634055 Tomsk, Russia*; C. CLAVEAU, J.-L. TEFFO, *Laboratoire de Physique Moléculaire pour l'Atmosphère et l'Astrophysique, CNRS UMR 7092, case 76, Université Pierre et Marie Curie, 4 Place Jussieu, F-75252 Paris Cedex 05, France*; J. VANDER AUWERA, *Service de Chimie Quantique et Photophysique C.P. 160/09, Université Libre de Bruxelles, 50 Avenue F. D. Roosevelt, B-1050 Brussels, Belgium*.

This work deals with the modeling of line intensities of cold and hot bands of  $^{13}\text{C}^{16}\text{O}_2$  corresponding to  $\Delta P = 5$  transitions ( $P = 2V_1 + V_2 + 3V_3$ ), using the effective operators approach.<sup>a</sup> In Brussels, 5 unapodized absorption spectra of  $^{13}\text{C}^{16}\text{O}_2$  (98% purity,  $P \times \ell = 7.43 - 538 \text{ mbar} \times \text{m}$ , room temperature) were recorded at a resolution of  $0.0035 \text{ cm}^{-1}$  (MOPD = 257.1 cm) using a Bruker IFS120HR. 382 absolute line intensities were measured in 9 bands observed between 3090 and 3920  $\text{cm}^{-1}$ . In Paris, 2 apodized absorption spectra of an  $^{18}\text{O}$ -enriched  $\text{CO}_2$  sample (297.0 K,  $\ell = 8.07(2)$  and  $36.07(4)$  m,  $P = 1.685(8)$  torr) containing 0.5% of  $^{13}\text{C}^{16}\text{O}_2$  were recorded at a resolution of  $0.0043 \text{ cm}^{-1}$  using a home-made FTS. 90 absolute line intensities were measured for 5 bands observed from 3400 to 3700  $\text{cm}^{-1}$ . A least squares fit of 479 observed line intensities, including those measured in this work and published for the 21102-00001 band,<sup>b</sup> allowed determination of 12 effective dipole moment parameters corresponding to the  $\Delta P = 5$  series of transitions (weighted standard deviation  $\chi = 0.855$ , RMS deviation = 4.69%). The new set of effective dipole moment parameters will soon be used to update the *Carbon Dioxide Spectroscopic Databank* (CDSD-1000).<sup>c</sup> Results will be presented and discussed.

<sup>a</sup>S.A. Tashkun, V.I. Perevalov, J.-L. Teffo, L.S. Rothman, and V.I.G. Tyuterev, *J. Quant. Spectrosc. Radiat. Transfer* **60**, 785–801 (1998); J.-L. Teffo, O.M. Lyulin, V.I. Perevalov, and E.I. Lobodenko, *J. Mol. Spectrosc.* **187**, 28–41 (1998).

<sup>b</sup>Y. Ding, P. Macko, D. Romanini, V.I. Perevalov, S.A. Tashkun, J.-L. Teffo, S. Hu, and A. Campargue, *J. Mol. Spectrosc.* **226**, 146–160 (2004).

<sup>c</sup>S.A. Tashkun, V.I. Perevalov, J.-L. Teffo, A.D. Bykov, and N.N. Lavrentieva, *J. Quant. Spectrosc. Radiat. Transfer* **82**, 165–196 (2003); <http://cdsd.iao.ru>, <http://cdsd.lpma.jussieu.fr>, <ftp://ftp.iao.ru/pub/CDSD-1000>.