

MILLIMETER WAVE SPECTROSCOPY AND MQDT CALCULATIONS OF HIGH RYDBERG STATES OF KRYPTON

MARTIN SCHÄFER and FRÉDÉRIC MERKT, *ETH Zürich, Laboratorium für Physikalische Chemie, ETH-Hönggerberg HCI, CH-8093 Zürich, Switzerland.*

A phase-stabilised backward wave oscillator (BWO) in the 260–380 GHz range was combined with a VUV laser system to record high-resolution spectra of high- n Rydberg states of krypton. Krypton atoms were excited into np ($n=58,60$) Rydberg states via the $4d[1/2]$ ($J = 1$) state using VUV and visible laser photons. Millimeter wave transitions between np and ns or nd Rydberg states were detected by pulsed field-ionization, at sub-MHz resolution and with mass selection. Using this excitation scheme, very accurate relative energies of fine and hyperfine structure levels of $(n + 2)s$ and nd ($n=68-74$) Rydberg states of ^{84}Kr and ^{83}Kr were obtained. In this region, s-d interactions are observable for ^{83}Kr due to the hyperfine interaction.

A multichannel quantum defect theory (MQDT) treatment of the hyperfine structure^a was used to analyze the millimeter wave data in combination with recent high-resolution VUV laser data^a and the available data from the literature; improved MQDT parameters and hyperfine structure data of the ^2P ground electronic state of $^{83}\text{Kr}^+$ were obtained.

^aH. J. WÖRNER, U. HOLLENSTEIN, AND F. MERKT, *Phys. Rev. A*, **68**, 032510 (2003).