

THE QUANTUM SOLVATION OF CYANOACETYLENE WITH HELIUM ATOMS: MICROWAVE SPECTROSCOPIC STUDIES

WENDY C. TOPIC and WOLFGANG JÄGER, *Department of Chemistry, University of Alberta, Edmonton, AB T6G 2G2, Canada.*

The rotational spectra of He_N-cyanoacetylene clusters ($N = 1, 2, 3, 4, 5..$) have been measured in the 3-27 GHz range using a high-resolution Fourier transform microwave spectrometer. The weakly bound clusters were generated in a pulsed free jet expansion. Manipulating the sample and nozzle conditions allowed some degree of control over the successive solvation of the molecule with He-atoms. Both the HCCCN and DCCCN isotopomers have been studied, allowing information about cluster geometry and dynamics to be determined. The presence of ¹⁴N and D nuclei in the clusters causes nuclear quadrupole hyperfine splitting of the He_N-HCCCN and He_N-DCCCN rotational transitions. This hyperfine structure was resolved, assigned, and analysed for low- J transitions. The value of the rotational constants decreases towards the helium nanodroplet value with increasing cluster size for small clusters. A "turn-around" in rotational constant value is expected for $N > 6$. Hyperfine fitting parameters and rotational constants will be interpreted in terms of the structure and dynamics of He_N-HCCCN clusters.