

## ROTATIONAL TRANSITIONS IN THE $\nu_9$ AND $\nu_7$ VIBRATIONAL STATES OF *cis*-HCOOH

OLEG I. BASKAKOV, *Department of Quantum Radiophysics, Kharkov National University, Svobody Sq. 4, 61077 Kharkov, Ukraine* ; BRENDA P. WINNEWISSER, IVAN R. MEDVEDEV, FRANK C. DE LUCIA, *Department of Physics, The Ohio State University, Columbus, OH 43210.*

The energy difference between *cis*-HCOOH and *trans*-HCOOH in the gas phase, ca.  $1365\text{ cm}^{-1}$ , was determined from a handful of MW transitions detected in formic acid at room temperature and reported in 1976<sup>a</sup>. A full analysis of the ground state spectrum of the *cis* rotamer based on FIR data was not provided until 2002<sup>b</sup>. For over 20 years unsuccessful efforts have been made to find vibrational bands of the *cis* rotamer in the gas phase IR spectrum. In the meantime, ab initio calculations and matrix spectra have yielded substantial information about the vibrational modes of this molecule<sup>c</sup>. We report here the first quantitative gas phase information about the excited vibrational states  $\nu_9$  and  $\nu_7$  of *cis*-HCOOH, obtained from FASSST measurements between 115 and 375 GHz, in a 6 m cell heated to 443 K. To date 427 transitions have been assigned for  $\nu_9$ , and 336 for  $\nu_7$ . The rotational and centrifugal distortion constants for these two states will be presented, together with a new set of ground state constants.

---

<sup>a</sup>W. H. Hocking, *Z. Naturforsch. A* **31**, 1113–1121 (1976).

<sup>b</sup>M. Winnewisser et al., *J. Mol. Spectrosc.* **216**, 259–265 (2002).

<sup>c</sup>E. M. S. Maçôas et al., *J. Mol. Spectrosc.* **219**, 70–80 (2003).