

INFRARED INTENSITY AS A SPECTRAL SIGNATURE OF H-BOND FORMATION IN SMALL AMMONIA CLUSTERS STUDIED IN HE DROPLETS

MIKHAIL N. SLIPCHENKO, *1605 Gilman Hall, Department of Chemistry, Iowa State University, Ames, IA 50011-3111*; BORIS G. SARTAKOV, *on leave from: General Physics Institute, Russian Academy of Sciences, Moscow, 119991, Russia*; ANDREY F. VILESOV, *Department of Chemistry, University of Southern California, Los Angeles, CA 90089, USA*.

Infrared spectra and intensities of the ν_1 , ν_3 H-stretching modes of $(\text{NH}_3)_n$ clusters ($n = 2-4$) have been obtained using the helium droplet isolation technique. Formation of a hydrogen bond in ammonia dimers leads to an increase of the infrared intensity by about a factor of four. In the larger clusters the infrared intensity per bond is close to that of dimers and approaches the value in the NH_3 crystal. Intensity of the overtone $2\nu_4$ band in trimers increases by a factor of 10 relative to that in single molecules and in dimers, and is comparable to the intensity of the ν_1 , ν_3 fundamental bands in larger clusters. These results provide new insights into the hydrogen bonding in ammonia clusters.