

PROGRESS ON THE INFRARED SPECTROSCOPY OF DIAMONDOIDS AND THEIR ASTROPHYSICAL APPLICATION

O. PIRALI, J. OOMENS, N. POLFER, *FOM Rijnhuizen, 3439MN Nieuwegein, The Netherlands*; Y. UENO, R. MABOUDIAN, *Department of Chemical Engineering, U.C. Berkeley, USA*; P. MAY, J. FILIK, *School of Chemistry, University of Bristol, UK*; J. DAHL, S. LIU, B. CARLSON, *Molecular Diamond Technologies, ChevronTexaco Technology Ventures, Richmond, USA*.

In the continuous effort for the unambiguous assignment of large carbonaceous species, we will present our recent laboratory observations concerning the infrared spectra of higher diamondoids (molecules consisting of diamond-like carbon cages) obtained by using various spectroscopic techniques. Increasing the size of the sample from adamantane ($C_{10}H_{16}$) to diamantane ($C_{14}H_{20}$), triamantane ($C_{18}H_{24}$), etc, corresponds to an addition of a cage C_4H_4 entity. Beyond its very interesting spectroscopic interest (due to the high symmetry of the systems studied here), this family of compounds which has been found in meteorites^a, also display a strong vibrational transition around 3.5 microns, thereby providing a convincing agreement with observed interstellar emission bands^b. We will discuss in this communication the spectroscopic analysis of our different spectra as well as our search for the possible occurrence of these compounds in several astrophysical observations.

^aE. Anders and E. Zinner, *Meteorites* 28 (1993) 490

^bL. J. Allamandola, S.A. Sandford, A.G. G. M. Tielens, T. M. Herbst, *Astrophysical Journal* 399 (1992) 134