

PULSED-FIELD IONIZATION ELECTRON SPECTROSCOPY OF COPPER-PYRAZINE, -PYRIMIDINE, AND -PYRIDAZINE COMPLEXES

XU WANG, JUNG SUP LEE, DONG-SHENG YANG, *Department of Chemistry, University of Kentucky, Lexington, KY 40605-0055.*

The copper complexes of pyrazine (1,4-C₄H₄N₂), pyrimidine (1,3-C₄H₄N₂), and pyridazine (1,2-C₄H₄N₂) are produced in laser-vaporization supersonic molecular beams and studied with pulsed field ionization zero electron kinetic energy spectroscopy and *ab initio* calculations. Adiabatic ionization energies of Cu-pyrazine, -pyrimidine and -pyridazine are measured to be 46038(7), 45332(7) and 43054(6) cm⁻¹, respectively. The metal-ligand stretching frequencies of Cu⁺/Cu-pyrazine, -pyrimidine, and -pyridazine are 265/190, 270/189, and 273/192 cm⁻¹, and the metal-ligand bending frequencies of these complexes are 115/74, 161/151, and 131/123 cm⁻¹, respectively. In addition, a symmetric ring stretch of 906 cm⁻¹ is observed for the Cu⁺-pyrazine ion. In all three complexes, Cu binds with one nitrogen atom and is located in the same plane as the ligand ring.