

ASSIGNMENT OF GG , GG' , TT and TG CONFORMERS IN THE *FASSST* ROTATIONAL SPECTRUM OF *N*-PROPANOL

Z. KISIEL, O. DOROSH, *Institute of Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warszawa, Poland*; A. MAEDA, F. C. DE LUCIA, E. HERBST, *Department of Physics, The Ohio State University, Columbus, OH 43210*.

Recent broadband measurements of the rotational spectrum of *n*-propanol up to 375 GHz resulted in determination of precise spectroscopic constants for the Gt conformer of this molecule.^a This is most likely the most stable conformer, although four other conformers are predicted to be very similar in energy. Assignment of some of these in cm-wave rotational spectra has previously been reported, but it was not possible to extend that work directly to mm-wave spectra.

Application of graphical Loomis-Wood techniques built into AABS^b and CAAARS^c spectral analysis packages eventually allowed successful assignment of the remaining four conformers: Gg , Gg' , Tt , and Tg *n*-propanol. It was realised that rotational energies in (Gg , Gg') and (Tt , Tg) pairs of conformers are highly coupled, but also amenable to description in terms of the Coriolis interaction mechanism. This allowed very precise determination of some energy level differences, such as $\Delta E(Gg' - Gg) = 3.035046(7) \text{ cm}^{-1}$. The assignment was checked against *ab initio* calculations, and is supported by new, precise determinations of dipole moments of some conformers, which were carried out using supersonic expansion cavity-FTMW spectroscopy.

^aA.Maeda, F.C.De Lucia, E.Herbst, J.C.Pearson, J.Riccobono, E.Trosell, R.K.Bohn, *Astrophys. J. Suppl.Series* **162**, 428 (2006)

^bZ.Kisiel, L.Pszczółkowski, I.R.Medvedev, M.Winnewisser, F.C.De Lucia, E.Herbst, *J. Mol. Spectrosc.* **233**, 231 (2005)

^cI.R.Medvedev, M.Winnewisser, B.P.Winnewisser, F.C.De Lucia, E.Herbst, *J. Mol. Struct.* **742**, 229 (2005)