

TUNABLE VACUUM ULTRAVIOLET PHOTOIONIZATION MEASUREMENTS OF CARBON CLUSTERS

B. W. TICKNOR, S. E. WHEELER, H. F. SCHAEFER III and M. A. DUNCAN, *Department of Chemistry, Univeristy of Georgia, Athens, GA, 30602-2556*; L. BELAU and M. AHMED, *Chemical Sciences Division, Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*; S. R. LEONE, *Departments of Chemistry and Physics and Ernest Orlando Lawrence Berkeley National Laboratory, Univeristy of California, Berkeley, California 94720*.

The photoionization thresholds for neutral carbon clusters have been measured using tunable vacuum ultraviolet light at the Advanced Light Source (ALS). A laser ablation cluster source with a high repetition rate vaporization laser is coupled to the quasi-continous light from the ALS. Ionized clusters are detected as a function of photon wavelength to determine the onset of ionization. Ionization thresholds for carbon clusters up to C₁₅ have been determined. These thresholds are compared to ionization potentials calculated from ab initio methods (extroplated CCSD(T) energies computed at cc-pVTZ ROCCSD(T) optimized geometries), which allows the investigation of isomeric structures in the molecular beam for some cluster sizes.