

## HIGH-RESOLUTION NEAR-INFRARED AND VISIBLE SPECTROSCOPY OF $\text{H}_3^+$

CHRISTOPHER P. MORONG, CHRISTOPHER F. NEESE AND TAKESHI OKA, *Department of Chemistry, Department of Astronomy & Astrophysics, and the Enrico Fermi Institute, University of Chicago, Chicago, IL 60637.*

Seven new rovibrational transitions of  $\text{H}_3^+$  have been observed in the visible region between 12,500-13,700  $\text{cm}^{-1}$  and at least eight new transitions have been observed in the near-infrared between 10,700-11,000  $\text{cm}^{-1}$ . These energy levels are above the barrier to linearity ( $>10,000 \text{ cm}^{-1}$ ), the regime in which  $\text{H}_3^+$  has enough energy to sample linear configurations. A high-resolution, high-sensitivity spectrometer based on a Ti:sapphire laser and incorporating velocity modulation and phase modulation with heterodyne detection<sup>a</sup>, was used to observe the transitions. The transitions are more than 6200 times weaker than the fundamental band. Due to the abundance of strong hydrogen Rydberg transitions, both pure hydrogen and He/ $\text{H}_2$  plasmas were used to identify the much weaker  $\text{H}_3^+$  transitions. The sparsity and weakness of the lines necessitated the use of the predicted intensities and frequencies<sup>b,c</sup> to focus our search. The measured rovibrational energy levels will assist in the development and verification of the theoretical calculations of  $\text{H}_3^+$  from first principles.

---

<sup>a</sup>J. Gottfried, B. McCall, and T. Oka, *J. Chem. Phys.* **118**, 10890 (2003).

<sup>b</sup>L. Neale, S. Miller, and J. Tennyson, *Astrophys. J.* **464**, 516 (1996).

<sup>c</sup>P. Schiffels, A. Aljiah, and J. Hinze, *Mol. Phys.* **101**, 189 (2003).