

IN-SITU PHOTOLYSIS OF METHYL IODIDE IN SOLID PARA-HYDROGEN AND SOLID ORTHO-DEUTERIUM

YUKI MIYAMOTO, *Division of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan*; MIZUHO FUSHITANI, *Institute für Experimentalphysik, Freie Universität Berlin, Arnimallee 14, 14195, Berlin, Germany*; HIROMICHI HOSHINA, *Terahertz Sensing and Imaging Laboratory, RIKEN; 519-1399 Aramaki Aoba, Sendai 980-0845, Japan*; and TAKAMASA MOMOSE, *Department of Chemistry, The University of British Columbia, Vancouver, B.C., V6T1Z1, Canada*.

Products of *in-situ* photolysis of methyl iodide isolated in solid para-hydrogen and solid ortho-deuterium were investigated by high-resolution FTIR spectroscopy. The ν_3 and ν_4 vibrational transitions of a methyl radical and the $^2P_{1/2} \leftarrow ^2P_{3/2}$ transition of an iodine atom were clearly observed in both solids after the photolysis, which indicates a small cage effect in hydrogen crystals. The vibrational transitions of the methyl radical showed a rotational structure subjected to crystal field splittings. Temperature dependence of the spectra allowed us to analyze the spectral fine structure completely based on our crystal field theory.^a On the other hand, the magnetic dipole transition of the iodine atom showed a rotational satellite induced by hydrogen molecules. The rotational satellite indicates a strong electron-rotor coupling in solid hydrogen. Details of the analysis will be discussed.

^aT. Momose, H. Hoshina, M. Fushitani, and H. Katsuki, *Vib. Spectrosc.* **34**, 95 (2004).