

SINGLE SHOT ULTRAFAST ELECTRON DIFFRACTION

F.M. RUDAKOV, J.D. CARDOZA, P.M. WEBER, *Department of Chemistry, Brown University, 324 Brook St, Providence, RI 02912, USA;*; D.H. DOWELL, J.F. SCHMERGE, J.B. HASTINGS, *Stanford Linear Accelerator, 2575 Sand Hill Rd, Menlo Park, CA 94025, USA.*

The time resolution of ultrafast electron diffraction experiment is primarily limited by the electron pulse duration. The radio frequency (rf)-guns are known to be capable of creating sub-picosecond pulses of relativistic electrons. We performed both experimental and computational studies of electron diffraction using the rf-gun at the Gun Test Facility located at the Stanford Linear Accelerator. The copper photocathode of the rf-gun is illuminated by the 1mm diameter beam from a quadrupled Nd:Glass laser beam at 263 nm, with a pulse width of 2 ps. The electrons scatter from a piece of 160 nm thick aluminum foil placed at 0.755 m from the photocathode, which subsequently get detected 4.7 m downstream. We observed the diffraction pattern obtained with a single electron pulse of 2 pC (10^7 electrons) and having a duration of 500 fs. Our simulations show that by reducing the laser pulse duration and the charge per pulse the time resolution of 100 fs is readily attainable.