

## RO-VIBRATIONAL ANALYSIS OF THE $\nu_4$ , $\nu_6$ and $\nu_3$ BANDS OF THIOFORMALDEHYDE: EXAMPLE OF A MASSIVE Z-TYPE CORIOLIS RESONANCE

W. J. LAFFERTY, *Optical Technology Division, NIST Gaithersburg, MD 20899, USA*; J.-M. FLAUD, A. PERRIN, *Laboratoire Inter Universitaire des Systemes Atmosphériques, CNRS, Universités Paris 12 et 7, 61 Av du Général de Gaulle, 94010 Créteil Cedex France*; H. BECKERS, Y.S. KIM, H. WILLNER, *Anorg. Chemistry, University of Wuppertal, D-42119 Wuppertal, GERMANY*.

The infrared spectrum of thioformaldehyde ( $\text{CH}_2\text{S}$ ) is of interest since this species is one of the myriad of molecules found in the interstellar space. From a spectroscopic point of view, it is of even more interest since the two lowest vibrational modes, the in-plane rocking mode,  $\nu_6$  ( $\text{B}_2$ ), and the out-of-plane wagging mode,  $\nu_4$  ( $\text{B}_1$ ), fall at 990.18 and 991.02  $\text{cm}^{-1}$  respectively. This separation of only 0.84  $\text{cm}^{-1}$  leads to a massive z-type Coriolis resonance where many of the rotational levels of each of the two vibrational states are mixed nearly 50% with each other. To make the situation even more interesting the C=S stretching vibration,  $\nu_3$ , with  $\text{A}_1$  symmetry occurs nearby at 1059  $\text{cm}^{-1}$ . This vibrational level also interacts with the two low frequency modes which complicates the assignment and analysis.  $\text{CH}_2\text{S}$  was produced by low pressure thermolysis of a gas flow of  $\text{C}_3\text{H}_5\text{SCH}_3/\text{Ar}$  (560° C) and  $\text{CH}_3\text{SCI}/\text{Ar}$  (1150° C) in the entrance of the multipath white cell (optical path length 32 m). At a total pressure of 0.15 mbar, 40 scans were recorded for the range 750 to 1400  $\text{cm}^{-1}$  on a Bruker HR120 TFIR spectrometer at a resolution of 0.005  $\text{cm}^{-1}$  (maximum optical path difference). The initial line assignment was not straightforward. There are strong series apparent in the spectrum, but the features expected for a b-type and c-type bands were not obvious near the band center. The centers of these three vibrations have been determined from medium resolution FT spectra<sup>a</sup> as well as laser Stark measurements<sup>b</sup>. An initial calculation was made using this information as well as guessed values for the band intensities. This permitted the identification of several low  $\text{K}_a$  series. Finally after numerous iterations, the transitions in the spectrum were identified leading to an excellent set of ro-vibrational constants.

<sup>a</sup>C. Clouthier, D.C. Moule, D.A. Ramsay and F.W. Birss, *Can. J. Phys* 60, 1212 (1982)

<sup>b</sup>D.J. Bedwell and G. Duxbury, *J. Mol. Spectrosc.* 64, 531 (1980)