

## INFRARED SPECTRUM OF MATRIX-ISOLATED METHYLHYDROXYCARBENE

C. A. BAUMANN, *Department of Chemistry, The University of Scranton, Scranton, PA 18510-4626.*

The ultraviolet irradiation of matrix-isolated biacetyl results in the formation of a complex of methylhydroxycarbene ( $\text{CH}_3\text{COH}$ ) and ketene ( $\text{CH}_2\text{CO}$ ). The infrared spectrum of the carbene exhibits broad features at  $3270\text{ cm}^{-1}$  (OH stretch) and  $1240\text{ cm}^{-1}$  (CO stretch); the vibrational frequencies for  $\text{CH}_2\text{CO}$  are shifted from those observed for the uncomplexed molecule. The complex was also observed when the parent was fully or partially deuterated, with appropriate isotopic shifting. The observed spectra are consistent with predictions from DFT and MP2 calculations for *trans*- $\text{CH}_3\text{COH}$  hydrogen bonded to the adjacent  $\text{CH}_2\text{CO}$ . The wavelength, matrix and isotope dependence for the formation of this complex was studied: parent deuteration decreases the apparent efficiency of production at 405 nm by a factor of 11 ( $\text{O}_2$  matrix) to 300 (Kr matrix).