

CO₂-BROADENED H₂O IN THE PURE ROTATION AND ν_2 FUNDAMENTAL REGIONS.

LINDA R. BROWN^a, *Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109*;
CATLIN M. HUMPHREY and ROBERT R. GAMACHE, *Department of Environmental, Earth & Atmospheric Sciences, University of Massachusetts School of Marine Sciences, 265 Riverside Street, Lowell, MA 01854-5045*.

The CO₂-broadened water coefficients (half-widths, line shifts and temperature dependence of the widths) are predicted using a fully complex Robert-Bonamy formulation for the 937 allowed and forbidden perpendicular type transitions of (000)-(000) between 200-900 cm⁻¹ in order to facilitate atmospheric remote sensing of the Mars and Venus. In addition, empirical Lorentz linewidths and pressure-induced frequency-shifts of CO₂-broadened H₂¹⁶O are obtained at room temperature for 257 perpendicular transitions of the (010)-(000) fundamental. For this, calibrated spectra recorded at 0.0054 cm⁻¹ resolution are measured assuming Voigt line shapes. For transitions between 1287 and 1988 cm⁻¹ with rotational quanta up to J = 13 and Ka = 6, the widths vary from 0.045 to 0.212 cm⁻¹ atm⁻¹ at 300 K; the pressure-shifts are quite large and range from -0.0386 to +0.0436 cm⁻¹ atm⁻¹. Observed and calculated values are compared.

^aSupport by NASA and NSF is gratefully acknowledged