THE ZEEMAN EFFECT IN THE OPTICAL SPECTRUM OF MANGANESE MONOHYDRIDE: MnH[a].

JAMIE GENGLER and TIMOTHY C. STEIMLE, *Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ, USA 85287*; MICHAEL STOLL, *Molecular Physics, Fritz-Haber-Institut der Max-Planck-Gesselschaft, Faradayweg 4-6, 14195 Berlin, Germany*.

The only successful demonstration of buffer-gas cooling and magnetic trapping of paramagnetic molecules was that performed by Doyle and co-workers in 1998[b]. It is anticipated that the relatively large rotational spacing and small spin-spin and spin-rotation interaction in the $X^7\Sigma^+$ (v = 0) state of MnH is a signature of a favorable ratio of elastic to inelastic collisions with helium atoms which is required for buffer-gas cooling and trapping. Furthermore, the large magnetic moment of the $N = 0$, $X^7\Sigma^+$ (v = 0) state of MnH enables one to create a deep trap, and this allows a large enough trapping time to be able to isolate the sample from its environment by cryopumping the helium away. Here we report on the magnetic properties of MnH studied as a necessary preliminary to our use of it in buffer-gas cooling and magnetic trapping experiments. The Zeeman effect in the $P_1(0)$ branch feature (ν = 17568.3536 cm^{-1}) of the $A^7\Pi - X^7\Sigma^+$ (0,0) band system of MnH was selected for Zeeman measurements. The field free spectra is complicated but has been thoroughly analyzed[c,d,e].

[a]Supported by the National Science Foundation-Exp. Phys. Chem.

[b]J.D. Weinstein, R. deCarvalho, T. Guillet, B. Friedrich, and J.M. Doyle, *Nature* **395**, 148 (1998).
[c]T.D. Varberg, R.W. Field, and A.J. Merer, *J. Chem. Phys.* **95**(3), 1563-1576 (1991).
[d]T.D. Varberg, J.A. Gray, R.W. Field, and A.J. Merer, *J. Mol. Spectrosc.* **156**, 296-318 (1992).
[e]J. Gengler, T.C. Steimle, J. Harrison, and J.M. Brown, *J. Mol. Spectrosc.* **241**, 192 (2007).