

LASER INDUCED FLUORESCENCE SPECTROSCOPY OF THE SiNSi ${}^2\Delta - \tilde{X}^2\Pi$ TRANSITION

MASARU FUKUSHIMA and TAKASHI ISHIWATA, *Faculty of Information Sciences, Hiroshima City University, Asa-Minami, Hiroshima 731-3194, Japan.*

On our SiCN/SiNC experiment^a, we measured laser induced fluorescence excitation spectrum in the range, 27,500 ~ 33,000 cm⁻¹, under jet cooled condition. Some of the vibronic bands in the spectrum have been assigned to Si containing species, such as Si₂, SiCH₂, SiCN, and SiNC, but some remain unassigned. The strongest band among them has a ${}^2\Pi - {}^2\Pi$ type rotational structure. The rotational constants of the ground and excited states are determined to be about 0.11 cm⁻¹, and they are about half of those of SiCN and SiNC. The spin-orbit interaction constant is determined to be 141.7 and 2.7 cm⁻¹ for the ground and excited states, respectively. Those are comparable with 140.8 and 4.9 cm⁻¹ for the $\tilde{X}^2\Pi$ and $\tilde{A}^2\Delta$ states of SiCN (not SiNC), respectively. As a result of some discussion, we have finally conclude that the spectrum can be attributed to the ${}^2\Delta - \tilde{X}^2\Pi$ transition of Si-N-Si. The ${}^2\Sigma^+ - \tilde{X}^2\Pi$ transition of Si₂N was studied by two groups^{b,c}. The rotational constant of the ground $\tilde{X}^2\Pi$ state agrees with that of the previous work. The vibronic structure of the laser induced fluorescence excitation spectrum is unusual; the ${}^2\Sigma - {}^2\Pi$ type vibronic bands have been only identified.

^aM. Fukushima and T. Ishiwata, *60th International Symposium on Molecular Spectroscopy*, TC02 (2005).

^bD. J. Brugh and M. D. Morse, *Chem. Phys. Lett.* **267**, 370 (1997).

^cZ. J. Jakubek, M. Zachwieja, S. G. Nakhate and B. Simard, *57th International Symposium on Molecular Spectroscopy*, MG05 (2002).